arXiv:1304.5727v1 [physics.plasm-ph] 21 Apr 2013

Ion-Wake induced anomaly of Dust Lattice Mode in presence of external Magnetic field

Saurav Bhattacharjee[1a)], Nilakshi Das[1b)]

[1] *Department of Physics, Tezpur University, Tezpur, Assam, 784 028, India*

Email: sauravtsk.bhattacharjee@gmail.com

we report a theoretical investigation of DL mode (Dust Lattice) in 2D Yukawa crystal in presence of asymmetric ion flow and external magnetic field perpendicular to the crystal plane. Two mutually perpendicular modes are found to be coupled due to Lorentz force. Interaction among the dust grains along vertical direction is strongly affected due to the formation of ion-wake. This causes anisotropy in interaction strength along two mutually perpendicular directions. Both hybrid modes are studied as characteristics of different ion flow speed and magnetic field strength. The study shows a fluctuation in DL mode frequency driven by the strength of the particle-wake interaction. Effect of ion flow on polarization of the hybrid wave amplitudes is discussed in details. Results show a possible mechanism of anomalous phase transition in dusty plasma.

Keywords: Dusty or complex plasmas, plasma dynamics and flow.

1. Introduction

The presence of micron sized solid particles in plasma adds a new dimension to it. Because of large and fluctuating charge on dust grains various novel and rich phenomena arise in plasma. Highly charged dust grains of nanometer to micrometer size interact with each other through repulsive Yukawa potential. Under the influence of strong repulsive potential, particles may form fluid and crystalline structures. This provides an interesting phase of dusty plasma that facilitates the study of various problems of fundamental physics like phase transition, crystal defects etc. Research in dusty plasma has observed an exponential growth after the discovery of plasma crystal by two pioneering groups led by Chu *et.al.*[1] and Thomas *et.al.*[2] in 1994. The crystal structures include, starting from the 1D chain, to 2D and 3D cubic crystals having different lattice geometries as Triangular, FCC, BCC and Hexagonal etc. Recent experiments in microgravity conditions provide an extended system of complex plasma for further investigations.

The discovery of low frequency dust mode in plasma [3] has triggered interest in studying other waves like dust ion-acoustic wave(DIA)[4] , electrostatic dust ion cyclotron waves (EDIC)[5], dust lattice waves (DLW) [6] . Melandsø[7] predicted theoretically the existence of dust lattice waves in 1996. This mode was experimentally observed in low temperature laboratory dusty plasma discharges by Homann *et.al.*[8] in 1997. In recent years, a variety of linear and nonlinear modes of the dust lattice waves have been investigated both experimentally and theoretically[9-10] . In presence of magnetic field perpendicular to crystal plane, two mutually perpendicular modes of a 2D yukawa crystal found to be coupled due to lorentz force [11]. This leads to two type of hybrid modes of high and low frequency branches instead of two independent modes.

As dust crystals are formed near the sheath region, the plasma flow dynamics near the sheath give rise to various attractive potentials such as shadowing and wake potential etc. in addition to the repulsive Yukawa type of interaction among dust grains. Closer to the lower electrode the velocity of the ion stream towards the negatively charged electrode may lead to instabilities like fluctuation in particle positions[12-13] and develop a new kind of potential named as wake potential. This instability leaves an impact on the dust lattice modes as well. It is identified as the possible cause for melting of dusty plasma crystal. Ivlev *et.al.*[14] have given an interesting theory on anisotropic dust lattice mode. They have also reported the decrease in frequency of the dust lattice modes due to particle wake interaction. Joyce *et.al.*[15] in 2001 for the first time presented a simulation model to study the behaviour of plasma crystal, incorporating both short range Yukawa potential as well as long range wake potential. They reported the formation of vertical strings of dust grains along the downward flow direction.

Ion-wake structure and their dynamics have been extensively studied in absence of magnetic field. In presence of magnetic field, very limited theoretical and experimental studies [16-17] are available. In the limit of linear response theory, Nambu *et.al.*[16] have reported damping of wake strength in presence of magnetic field. This leads to a weak effective interaction strength between dust grains along vertical direction as compared to unmagnetized case and is verified experimentally[17] in recent year. In a recent paper Saurav *et.al.* [18]

have investigated the effect of wake potential on Coulomb crystallization in presence of magnetic field. They observed a reduction in RDF (Radial Distribution Function) peak heights with increased strength of wake potential and reported a ion-wake induced dislocation of dust grains from regular crystal arrangement. This modification in particle-wake interaction in presence of magnetic field with the strength of the ion flow and its effect on various eigen modes is an important issue in complex plasma.

The phase behaviour of fluids depends solely on the intermolecular interaction. Understanding this mechanism is one of the biggest challenges in statistical thermodynamics. Ion-wake formation and its dynamics found to play a vital role in understanding the formation of coulomb crystals and their phase transition in magnetized complex plasma. In this letter we presented a theoretical study to investigate the behaviour of Dust lattice modes in presence of magnetic field perpendicular to crystal plane and ion flow. Polarization of both the hybrid wave amplitudes and their interplay with the magnetic field have been discussed in details. Results show a possible kind of anomalous phase transition[19] in dusty plasma.

2. Theoretical Formulation

2.1. Interaction Potential among dust grains

We consider a 2D uniform dusty plasma with stationary heavier dust grains in presence of an external magntetic field B_o $\widehat{x}$ perpendicular to the downstream ion flow. Under the influence of magnetic field, the ion flow dynamics get affected and subsequently the dielectric response function [16] of the plasma medium also changes. In presence of low frequency wave ($\omega \ll \omega_{ce}$),with phase velocity smaller than electron thermal velocity, electrons will have a Boltzmanian distribution. In normalized scale of electron Debye length λ_{De}, the dispersion relation can be written as:

$$\frac{1}{k^2\epsilon(\omega,k)} = \frac{1}{k^2+1} + \frac{{k_\parallel}^2}{(k^2+1)M^2({k_\parallel}^2-{k_R}^2)({k_\parallel}^2-{k_{Im}}^2)} \tag{1}$$

where it is assumed that $k_\parallel \gg k_y$ and $k^2 = {k_z}^2 + {k_y}^2$, $k_z(\parallel)$ and $k_y(\perp)$ are wave vectors parallel and perpendicular to the direction of ion flow. Here M=$\frac{u_{io}}{C_i}$, is the mach number having $c_i = \lambda_{De}\omega_{pi}$ as ion acoustic wave velocity and u_{io} is the streaming velocity. $f = \frac{{\omega_{pi}}^2}{{\omega_{ci}}^2}$, ω_{pi} and ω_{ci} are the ion plasma frequency and ion cyclotron frequency respectively. k_R and k_{Im} corresponds to real and imaginary screening modes as a function of $k_\perp$, in a complex $k_\parallel$ plane.

$${k_{R,Im}}^2 = -\frac{1}{2}({k_\perp}^2 + 1 - \frac{1}{fM^2} - \frac{1}{M^2}) \pm \sqrt{\frac{1}{4}({k_\perp}^2 + 1 - \frac{1}{fM^2} - \frac{1}{M^2})^2 + \frac{{k_\perp}^2+1}{M^2 f}} \tag{2}$$

In the limit of $k_\perp > \frac{1}{fM^2}$, magnetic field B<1.0T, for all M, both the pole can be approximated as $k_R \approx \sqrt{\frac{1}{fM^2}}$, $k_{Im} \approx \pm i\sqrt{{k_\perp}^2 + 1 - \frac{1}{M^2}}$.

Using the dispersion relation in equation (1) one can write the interaction potential between the test dust particulates in presence of magnetic field as

$$\Phi(y,z) = \frac{Q_d}{(2\pi)^3\lambda_{De}\epsilon_0}\int d^3k\delta(k)[\frac{1}{k^2+1} + \frac{{k_\parallel}^2}{(k^2+1)M^2({k_\parallel}^2-{k_R}^2)({k_\parallel}^2-{k_{Im}}^2)}]e^{ik.r} \tag{3}$$

where r=$\sqrt{y^2+z^2}$. Integrating the first part on the R.H.S of the above equation gives Yukawa potential

$$\Phi_D = \frac{Q_d}{4\pi\epsilon_0 r}exp(\frac{-r}{\lambda_{De}}) \tag{4}$$

Integrating the second part on R.H.S of equation (3) for the real pole gives oscillatory wake potential

$$\Phi_w(0,z) = -\frac{Q_d}{4\pi\lambda_{De}\epsilon_0}2\pi\frac{\sqrt{1/f}Msin(\frac{\sqrt{1/f}z}{M})}{(\frac{1}{f}+M^2)(\frac{1}{f}-1+M^2)} \tag{5}$$

In the limit of $f \gg 1$, above equation shows a damping in the strength of the wake potential.

2.2. Dispersion relation of Dust Lattice Wave (DLW)

We have considered a simplified model of monolayer hexagonal lattice in Y-Z plane with ion flow along downward z-direction and magnetic field perpendicular to the crystal plane. The lattice mode propagate along Y-direction parallel to the fundamental lattice vector directed from center of the cell to one of its nearest neighbors. The equation of motion along two transverse directions are given by [11]

$$m_d\frac{d^2y_{n1}}{dt^2} = \sum_{m=1,m\neq n}^{N} K_{y_{nm}}(y_{n1}-y_{m1}) + Q_dB_o\frac{dz_{n1}}{dt} \tag{6}$$

$$m_d\frac{d^2z_{n1}}{dt^2} = \sum_{m=1,n\neq m}^{N} [K_{z_{nm}}+K_{zw_{nm}}](z_{n1}-z_{m1}) - Q_dB_o\frac{dy_{n1}}{dt} \tag{7}$$

Where $y_{(n,m)1}$ and $z_{(n,m)1}$ are the displacement of the particles from their equilibrium positions. $K_{(y,z)_{nm}}$ are the spring constants corresponding to Yukawa type of interaction along y and z direction and $K_{zw_{nm}}$ is the spring constant corresponding to wake potential along vertical z direction.

Using equation (4) we can write the normalized spring constants $K_{y,z}$ corresponding to Yukawa potential along y and z direction as

$$K_{y,z} = \frac{d^2\Phi_D}{d(y,z)^2}|_{y,z=a} = \frac{\Gamma exp(-\sqrt{2}\kappa)}{2^{5/2}\kappa^2}[3+3\sqrt{2}\kappa+2\kappa^2] \tag{8}$$

Here a is average interparticle distance, Γ is coulomb coupling parameter and $\kappa = \frac{a}{\lambda_{De}}$ stands for screening constant.

Similarly from equation (5)we get the spring constant for the wake potential as

$$K_{zw} = 2\pi \frac{(1/f)^{3/2} sin(\frac{\sqrt{1/f}\kappa}{M})}{M(\frac{1}{f} + M^2)(\frac{1}{f} + M^2 - 1)} \tag{9}$$

This leads to the effective spring constant along z-direction as

$$K_z = K_z(Yukawa) + K_{zw}(Wake) \tag{10}$$

As we have considered that DL mode propagates along Y direction, the oscillations of the wave modes are y_{n1}=$A_y exp[i(k_y a - \omega t)]$ and z_{n1}=$A_z exp[i(k_y a - \omega t)]$ respectively along two mutually perpendicular directions. Substituting y_{n1} and z_{n1} in equation (6) and (7) we get the dispersion relation as:

$$[\omega^2 + \sum_{m=1,n\neq m}^{N} K_{y_{nm}}(\exp[ik_y a] - 1)][\omega^2 + \sum_{m=1,n\neq m}^{N} (K_{z_{nm}} + K_{zw_m})(\exp[ik_y a] - 1)] - {\omega_c}^2\omega^2 = 0 \tag{11}$$

The dispersion relation indicates that two transverse modes get coupled by Lorentz force and there exist two hybrid modes instead of independent modes.

In the limit of $k_y a \ll 1$, $\cos(k_y a) \approx 1 - \frac{1}{2}(k_y a)^2 + ...$ equation (11) gives two roots as

$$\omega_H^2 \approx \omega_c^2 + \sum_{m=1,n\neq m}^{N} 2(K_{y_{nm}} + K_{z_{nm}} + K_{zw_m}) \sin^2(\frac{k_y a}{2}), \text{[Upper Hybrid]} \tag{12}$$

$$\omega_L^2 \approx \frac{\sum_{m=1,n\neq m}^{N}[2K_{y_{nm}} \sin^2(\frac{k_y a}{2})] \sum_{m=1,n\neq m}^{N}[2(K_{z_{nm}} + K_{zw_m}) \sin^2(\frac{k_y a}{2})]}{\omega_c^2}, \text{[Lower Hybrid]} \tag{13}$$

Here the frequency is normalized by $\omega_N = \sqrt{\frac{K_B T_d}{m_d \lambda_{De}^2}}$, where K_B, T_d, m_d and λ_d be the Boltzmann constant, dust temperature, dust mass and dust Debye length respectively.

Substituting the displacements y_{nm} and z_{nm} in equation (6) and (7) we can get the wave amplitude ratio along y and z direction as:

$$\frac{y_{nm}}{z_{nm}} = i\mu, \text{where} |\mu| = \sqrt{\frac{\omega^2 + 2(K_z + K_{zw})(\cos(k_y a) - 1)}{\omega^2 + 2K_y(\cos(k_y a) - 1)}} \tag{14}$$

Using equation (14) one can write the polarization P of the wave amplitudes as

$$P = \frac{1}{1 + \frac{1}{|\mu|}} \tag{15}$$

P=1 corresponds to pure longitudinal mode and P=0 corresponds to pure transverse mode.

3. Results and Discussion

The Debye cloud surrounding the dust particles gets distorted by the downward ion flow and becomes elongated in lower vertical (z-direction) direction. Amount of distortion and the scale length of elongation depends mainly on the strength of the magnetic field and flow velocity of the downward ion stream towards the plasma sheath. As there is no attractive type of potential in the horizontal direction, this excess positive charge on the lower side of the Debye cloud attracts the dust particles just below it to align along vertical direction. This process of alignment has a vital dependency on the ion-wake induced vertical confinement frequency. It is obvious from equation (5) that frequency of the wake potential increases with decrease in Mach number for a given magnetic field strength. As dust particles are confined at the trail of the ion-wake, alignment continues up to a certain value of confinement frequency and beyond that dust particles are no more confined and they restore their original configuration once again, as demonstrated in Fig.1(a) . Fig. 1(b) clearly shows the transition of vertical confinement strength with ion flow speed.

FIG.1:(color online) (a) Vertical confinement mechanism of particle-wake interaction with increase in strength of wake potential from left to right. (b) Effective spring constant K_z vs. M(mach number).

Due to this interaction between dust particles and ion-wake, partciles are first dislocated from their regular orientation and then again restore their original configuration. This entire process leaves a strong impact on the dust lattice modes. It is visible from the dispersion relations (12) and (13), shown in Fig. 2, for different magnetic field strength and ion flow velocity.

FIG. 2: (color online) The plots show the dispersion relation profiles with different Mach number M and magnetic field strength B for coulomb coupling parameter Γ=824 and screening strength κ=2.9. (a) Shows the normalized frequency dependence of both Upper Hybrid (UH) (solid line) and Lower Hybrid (dotted line) waves with M=0.3(green), 0.18(blue) and 0.11(red), for a given magnetic field B=0.08T. (b) M=0.6 (green), 0.4(blue) and 0.25(red) with B=0.06T. (c) M=0.5 (green), 0.3 (blue) and 0.15(red) with B=0.04T.

It is evident from equations (14) and (15) that polarization of the hybrid wave amplitudes varies with ion flow speed as well as the magnetic field strength. At comparatively higher ion flow speed (M>0.4), the strength of the wake potential is not sufficient enough to affect the polarization of the wave amplitudes as can be seen from FIG 3(a). Above discussion clears the fact that effective repulsive strength along vertical direction first decreases with decrease in M value (strong confinement zone) due to enhancement in strength of the wake potential. In that region we find the lower hybrid mode becomes more longitudinal in nature as compared to upper hybrid for a given magnetic field strength and wave number.

FIG. 3: (color online) The profiles show (a) Polarization P vs. Mach number M, for both upper hybrid (UH) (green solid) and lower hybrid (LH) (green dotted) waves for a given wave number ka/π=0.6 and magnetic field strength B=0.08T. (b) P vs. M for a given wave number ka/π=0.6 and different magnetic fields B=0.08T, UH (green solid), LH (green dotted). B=0.06T, UH (blue solid), LH (blue dotted). (c) Strength of spring constant k_z vs. $f(B)={\omega_{pi}}^2/{\omega_{ci}}^2$ for a given plasma frequency ω_{pi}=1.86×10^7 s^{-1} and Mach Number M=0.5.

On further decrease in Mach number (weak confinement zone) below M$<$ 0.24, upper hybrid becomes more longitudinal in nature as compared to lower hybrid, shown in Fig.3(a). It can be seen from Fig. 3(c) that effective spring constants along vertical direction decreases with increase in strength of the magnetic field (decrease in f value) and agrees with the experimental result[17]. This causes shifting of the critical Mach number, from which confinement starts dominating, to a lower value, with decrease in strength of the magnetic field as shown in FIG. 3(b). This ion-wake induced periodic transition in crystal phase from a ordered to disorderd state and again get back to the ordered state showing an analogy with the anomalous phase transitions in complex fluids.

4. Conclusions

In the present piece of work we have elaborately discussed the effect of wake potential on dust lattice modes in terms of ion flow speed and external magnetic field for a 2D crystal structure. The observations can be summarized as follows:

(a) The ion flow near the sheath region gives rise to attractive wake potential along vertical direction and this causes anisotropy in the interaction strength along two mutually perpendicular directions.

(b) It is demonstrated in FIG. 1 that with increase in strength of the wake potential, the dust particles start aligning along vertical direction from their regular orientations in the crystal plane. On further increase in strength, confinement stops and particles regain their original configuration. Impact of this mechanism on DL modes is clear from dispersion relations in Fig.2.

(c) Polarization of both the hybrid wave amplitudes is also affected by the strength of the wake potential. Within the strong confinement zone upper hybrid mode is found to be more transversely polarized as compared to lower hybrid for a given magnetic field and wave number. On further decrease in M value, in the weak confinement zone of interaction, polarization of both the hybrid wave amplitudes reverses in character and upper hybrid mode becomes more longitudinally polarized as compared to lower hybrid. It can be seen from FIG. 3(a).

(d) With increase in strength of the magnetic field, effective interaction along z-increases. As a result, the critical Mach number is shifted to a lower value with decrease in strength of the magnetic field for a given wave number. It is shown in FIG. 3(b).

(e) Finally, we can say the particle-wake interaction is responsible for decrease in frequencies of the DL modes up to certain strength of wake potential. Beyond that DL mode frequency increases again with the strength of the wake potential.

5. Referrence

[1] J.H. Chu and I.Lin, Phys.Rev.Lett. 72, 4009 (1994).

[2] H. Thomas, G.E. Morfill, V. Demmel, J. Goree, B. Feuerbacher and D. Möhlmann, Phys. Rev. Lett. 73, 652 (1994).

[3] N.N. Rao, P.K. Shukla and M.Y. Yu, Planet. SpaceSci. 38, 4 (1990).

[4] P. K. Shukla and V. P. Silin, Physica Scripta. 45, 508 (1992).

[5] R. L. Merlino, A. Barkan, C. Thompson, and N. DAngelo, Phys.Plasmas, 5, 1607 (1998).

[6] X. Wang, A. Bhattacharjee and S. Hu, Phys.Rev.Lett. 86, 2569 (2001).

[7] F. Melandsø, Phys.Plasmas. 3, 3890 (1996).

[8] A. Homann, A. Melzer, S. Peters, and A. Piel, Phys. Rev. E 56, 7138 (1997).

[9] S. Nunomura, D. Samsonov and J. Goree, Phys.Rev.Lett. 84, 5141 (2000).

[10] T. Misawa, N. Ohno, K. Asano1, M. Sawai, S. Takamura, and P. K. Kaw, Phys. Rev. Lett. 86, 1219 (2001).

[11] G. Uchida, U. Konopka and G. Morfill, Phys.Rev.Lett. 93, 155002 (2004).

[12] A. Melzer, V.S. Schweigert and A. Piel, Physica Scripta 61, 494 (2000).

[13]V. Steinberg, R. Stterlin, A. V. Ivlev, and G. Morll, Phys.Rev.Lett. 86, 4540 (2001).

[14] A. V. Ivlev and G. Morfill, Phys. Rev. E 63, 016409 (2000).

[15] G. Joyce, M. Lampe and G. Ganguli, IEEE Transaction on Plasma Science, 29, 238 (2001).

[16] M.Nambu, M.Salimullah and R.Bingham, Phys.Rev.E 63, 056403 (2001).

[17] J. Carstensen, F. Greiner and A. Piel, Phys.Rev.Lett. 109, 135001 (2012).

[18] S.Bhattacharjee and N.Das, Phys.Plasmas 19, 103707 (2012).

[19] A . B . de Oliveira , E . B . Neves , C . G. Paukowski , J. Z . Netz, P. A. and M . C . Barbosa , J.Chem.Phys. 132, 164505 (2010).